


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 401/12, A61K 31/47, C07D 215/14, 215/18		A1	(11) International Publication Number: WO 96/04267
			(43) International Publication Date: 15 February 1996 (15.02.96)
(21) International Application Number: PCT/EP95/02970		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).	
(22) International Filing Date: 27 July 1995 (27.07.95)		Published With international search report.	
(30) Priority Data: P 9401696 1 August 1994 (01.08.94) ES			
(71) Applicant (for all designated States except US): LABORATORIOS MENARINI S.A. [ES/ES]; Alfonso XII, 587, E-08912 Badalona (ES).			
(72) Inventors; and (75) Inventors/Applicants (for US only): MAULEON CASELLAS, David [ES/ES]; Alfonso XII, 587, E-08912 Badalona (ES). CARGANICO, Germano [IT/ES]; Alfonso XII, 587, E-08912 Badalona (ES). FOS TORRO, María de los Desamparados [ES/ES]; Alfonso XII, 587, E-08912 Badalona (ES). GARCIA PEREZ, María Luisa [ES/ES]; Alfonso XII, 587, E-08912 Badalona (ES). PALOMER BENET, Albert [ES/ES]; Alfonso XII, 587, E-08912 Badalona (ES).			
(74) Agent: MINOJA, Fabrizio; Studio Consulenza Brevettuale, Via Rossini, 8, I-20122 Milano (IT).			

(54) Title: NAPHTHALENE AMIDES HAVING LEUKOTRIENE-ANTAGONISTIC ACTION



(57) Abstract

Naphthalene amides of formula (I) wherein the substituent containing A is bound to the 6- or 7- position of the 2-naphthol system; the substituent containing B is bound to the benzene ring at any free position; R¹ is hydrogen or methyl; R² is hydrogen, fluorine, chlorine or -OCH₃, which is bound to the naphthalene system at any positions except the 2- and the one occupied by the other substituent; R³ is hydrogen, fluorine, chlorine or bromine; A- is -CO-NR⁴- or -NR⁴-CO- group, wherein R⁴ is hydrogen or methyl; B is a 5-tetrazolyl or -COOR⁵ group, wherein R⁵ is hydrogen, a (C₁-C₄)-alkyl or a phenylalkyl group of less than 10 carbon atoms; m is 0 or 1; n and p are integers from 0 to 6, with the proviso that n + p is less or equal to 6; as well as the solvates and pharmaceutically acceptable salts thereof, have leukotriene antagonistic action.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

NAPHTHALENE AMIDES HAVING LEUKOTRIENE-ANTAGONISTIC ACTION

The present invention relates to novel naphthalene amides, the pharmaceutically acceptable salts and solvates thereof and pharmaceutical compositions containing them, having a leukotriene-antagonistic activity. The present invention also relates to a process for the preparation of the novel naphthalene amides, as well as to the therapeutic use thereof.

TECHNOLOGICAL BACKGROUND

It is well known that most eicosanoids, prostaglandins, leukotrienes and related compounds derive from a fatty acid having 20 carbons and 4 unsaturations, called arachidonic acid (AA), which fundamentally esterifies the hydroxyl at the 2- position of the glycerol of the phospholipids contained in the cell membranes. AA is released from the phospholipid containing it by the action of a lipase, phospholipase A₂ (PLA₂) ("CRC Handbook of Eicosanoids and Related Lipids", vol. II, Ed. A.L. Willis, CRS Press Inc., Florida (1989)). After being released AA is metabolized in mammals mainly by two different pathways or enzyme systems. Through cyclooxygenase it produces prostaglandins and thromboxanes, the most significant being PGE₂ and TxA₂, which are directly involved in inflammation (Higgs et al. Annals of Clinical Research, 16, 287 (1984)). Through lipo-oxygenase it produces leukotrienes, the most important being LTB₄, and the peptide-leukotrienes LTC₄, LTD₄ and LTE₄. All of them are also involved in inflammatory reactions, exhibiting

chemotactic activities, stimulating the secretion of lysosomic enzymes and playing an important role in immediate hypersensitivity reactions (Bailey and Casey, Ann. Rep. Med. Chem., 17, 203 (1982)). Leukotriene LTB₄ is a strong chemotactic agent which promotes the infiltration of leukocytes and their subsequent degranulation. (Salmon et al., Prog. Drug Res., 37, 9 (1991)). It has been widely shown that LTC₄ and LTD₄ have strong constrictive action on human bronchi (Dahlen et al., Nature, 288, 484 (1980)), causing the obstruction of airways by inflammation and mucus production (Marom et al., Am. Rev. Resp. Dis., 126, 449 (1982)), being thus involved in the pathogenesis of bronchial asthma, chronic bronchitis, allergic rhinitis, etc. Peptide-leukotrienes also bring about a blood extravasation caused by the increase of vascular permeability (Camp et al., Br. J. Pharmacol., 80, 497 (1983)) and are involved in some inflammatory diseases such as atopic eczema and psoriasis. On the other hand, several effects of peptide-leukotrienes on human cardiovascular system have been observed; they are mainly involved in the pathogenesis of the ischaemic cardiopathy. This relationship has been confirmed by the fact that coronary arteries can produce these mediators (Piomelli et al., J. Clin. Res., 33, 521A (1985)). These effects, together with the strong contractions observed in heart tissue caused by LTC₄ and LTD₄, suggest that these mediators might contribute to other cardiovascular disorders, such as coronary spasm, heart anaphylaxis, cerebral oedema and endotoxic shock.

From what said above it follows that the control of

the biological activity of leukotrienes through compounds which inhibit their release or antagonize their effects, represents a new rational approach to the prevention, elimination or improvement of different
5 allergic, anaphylactic, inflammatory and thrombotic conditions, in which such mediators are involved.


In literature some compounds have been described that can be considered as structurally related to the compounds of the present invention, having an inhibiting
10 action on enzyme 5-lypoxygenase, and a leukotriene antagonistic activity. Kreft A.F. et al. described 2-[6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid (US 4,960,892) and derivatives thereof (US 5,084,575), among which there are compounds containing sulfonimide,
15 hydroxamic acid or hydroxamate groups, but the amides of the present invention are not included in their general formulas.

On the other hand, Stevenson D. et al. (US 4,579,866) described amides having an inhibiting action
20 on 5-lypoxygenase, but differing from the compounds of the present invention in two aspects: first, they contain a phenyl instead of a naphthyl; second, they contain an alkyl chain with only one carbon atom with or without branches between the amide function and the
25 terminal carboxylic acid. In short, they are compounds containing a phenylalanine or glycine terminal residue.

The obtaining of compounds with high leukotriene antagonistic activity is still a problem in the therapy. The present invention provides a number of novel
30 compounds that exhibit the above mentioned antagonistic action and that are useful in therapy.

DISCLOSURE OF THE INVENTION

The present invention relates to novel naphthalene amides of general formula I,



10 wherein:

the substituent containing A is bound to the 6- or 7-position of the 2-naphthol system;

the substituent containing B is bound to the benzene ring at any free position;

15 -R¹ is hydrogen or methyl;

-R² is hydrogen, fluorine, chlorine or -OCH₃, which is bound to the naphthalene system at any position except the 2- and the one occupied by the other substituent;

-R³ is hydrogen, fluorine, chlorine or bromine;

20 -A- is a -CO-NR⁴- or -NR⁴-CO- group, wherein R⁴ is hydrogen or methyl;

-B is a 5-tetrazolyl or -COOR⁵ group, wherein R⁵ is hydrogen, a (C₁-C₄)-alkyl or a phenylalkyl group of less than 10 carbon atoms;

25 m is 0 or 1;

n and p are integers from 0 to 6, with the proviso that n + p is less or equal to 6.


The present invention also relates to a process for the preparation of the novel naphthalene amides, as well as the therapeutical use thereof.

30

The present invention also relates to the solvates

and the pharmaceutically acceptable salts of the amides of formula I and particularly the salts represented by formula Ia,

5



Ia

10 wherein M^+ is an alkali metal cation (e.g. Na^+ , K^+) or represents a half amount of an alkaline-earth metal cation (e.g. $1/2 Ca^{2+}$, $1/2 Mg^{2+}$), or a cation deriving from an amine or quaternary ammonium salt (such as triethanolammonium, tris(hydroxymethyl)methylammonium).

15 The compounds of formula I can have one or more asymmetric carbons in their structure. The present invention comprises all the possible stereoisomers as well as the mixtures thereof.

Preferred compounds are those wherein R^2 is hydrogen and B is a 5-tetrazolyl or $COOR^5$ group, wherein R^5 is hydrogen, methyl, ethyl or benzyl.

Also preferred are the compounds of formula I wherein R^3 is hydrogen or chlorine and -A- is -CONH- or -NHCO-.

25 When the substituent containing A is bound to the 6- position of the 2-naphthol system, particularly preferred are the compounds of formula I wherein R^1 is hydrogen, m is 1, and -A- is -NHCO-; or those wherein -A- is -CONH-, being n and p integers ranging between 0 and 3.

30 When the substituent containing a A is bound to the

7- position of the 2-naphthol system, particularly preferred are the compounds of formula I wherein R¹ is hydrogen, m is 1 and -A- is -CONH-; or also those wherein m is 0, -A- is -CONH-, being n and p integers ranging between 0 and 3.

Particularly preferred compounds of the present invention are the following ones: N-[4-(1H-5-tetrazolyl)phenylmethyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide;

10 N-[3-(1H-5-tetrazolyl)phenylmethyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide;

N-[2-(1H-5-tetrazolyl)phenylmethyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide;

15 N-[4-(1H-5-tetrazolyl)methylphenyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide;

N-[4-(1H-5-tetrazolyl)methylphenyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide (sodium salt);

4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamido]-benzoic acid;

20 4-[4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamido]phenyl]butanoic acid;

4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidomethyl]benzoic acid;

25 3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidomethyl]benzoic acid;

4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamido]-phenylacetic acid;


3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamido]-phenylacetic acid;

30 4-[4-[2-[6-[(7-chloro-2-quinolinyl)methoxy]-2-naphthyl]propanamido]phenyl]butanoic acid;


- N-[4-(1H-5-tetrazolyl)phenylmethyl]-6-(2-quinolinylmethoxy)-2-naphthaleneacetamide;
 4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]carboxamido]-phenyl]butanoic acid;
 5 N-[3-(1H-5-tetrazolyl)phenylmethyl]-6-(2-quinolinylmethoxy)-2-naphthalenecarboxamide;
 4-[4-[[7-(2-quinolinylmethoxy)-2-naphthyl]carboxamido]-phenyl]butanoic acid;
 N-[4-(1H-5-tetrazolyl)phenylmethyl]-7-(2-quinolinylmethoxy)-2-naphthalenecarboxamide;
 10 4-[2-[[7-(2-quinolinylmethoxy)-2-naphthyl]carboxamido]-ethyl]benzoic acid;
 N-[4-(1H-5-tetrazolyl)phenylethyl]-7-(2-quinolinylmethoxy)-2-naphthalenecarboxamide;
 15 4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]methylamino-carbonyl]phenyl]butanoic acid;
 N-[4-(1H-5-tetrazolyl)phenylpropyl]-7-(2-quinolinylmethoxy)-2-naphthalenecarboxamide;
 as well as the carboxylic acid esters described in the
 20 examples.

According to the present invention, the compounds of general formula I wherein A is $-\text{CO}-\text{NR}^4-$ are obtained by a process in which, starting from a compound of general formula II,

25



- 30 wherein R^1 , R^2 , R^3 and m have the above defined meanings, is reacted with a compound III,




5

III

wherein R^4 , n and p have the above defined meanings and D can be equivalent to the group B in I or, when B in formula I is COOH , then D contains a suitable carboxy-protecting group, for example as a methyl, ethyl or benzyl ester. The reaction between II and III is carried out in the presence of a carboxy-activating agent such as dicyclohexylcarbodiimide or 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide and a base such as triethylamine or 4-dimethylaminopyridine, in a suitable aprotic solvent such as chloroform, methylene chloride or N,N -dimethylformamide, at a temperature ranging between 0° and 40°C for a time from 3 to 24 hours. In this way, a compound of formula IVa is obtained,

20



25


IVa

this compound coincides with I or is converted into I removing any COOH -protecting groups present in D , then, when D is for example a methyl, ethyl or benzyl ester, can be removed by treatment with a suitable base such as lithium or sodium hydroxide in aqueous solution, in a suitable organic solvent such as methanol, ethanol or


30

tetrahydrofuran, at a temperature ranging between 20°C and the solvent reflux, for a time from 1 to 48 hours.

A compound of general formula I wherein A is -NR⁴-CO- is obtained according to the same process as above, starting from the compounds V and VI,




V



VI

wherein R¹, R², R³, R⁴, D, m, n and p have the above defined meanings. In this way a compound of formula IVb,



IVb

which coincides to I or is converted into I removing any COOH-protecting groups present in D, is obtained as described above.


When a given salt of general formula Ia is desired, a compound I can be treated with a suitable base or ion-exchanger, according to the conventional chemical techniques. Thus, for example, I can be treated with sodium hydroxide or tris(hydroxymethyl)methylamine in a suitable solvent such as water-methanol or ethanol mixtures, for a time from 15 min to 2 hours, at a

10

temperature ranging between 25°C and the solvent reflux.

A starting product of formula II can be obtained, for example, following the synthesis sequence shown in scheme 1.

Scheme 1




In this synthetic sequence, a compound IX can be obtained, for example, subjecting a compound VII to the action of a base such as sodium methoxide or sodium hydride, thereafter reacting it with a compound VIII, wherein R^3 represents the groups defined above and X is a bromine or chlorine atom, in a suitable organic solvent such as benzene, N,N-dimethylformamide or tetrahydrofuran, at a temperature ranging between 0° and 25°C for a time from 3 to 24 hours (step (1)).

A compound II can be obtained starting from IX (step (2)) by basic hydrolysis as described for the preparation of I with $B=COOH$ starting from IVa.

11

Analogously, a starting compound of general formula V can be obtained, for example, following the synthetic sequence shown in scheme 2.

Scheme 2




In this synthetic sequence a compound XI is obtained starting from a compound X (step (3)), following the same process as described for the preparation of IX.

A compound V can be obtained starting from XI by hydrolysis with hydrochloric or sulfuric acid, in a suitable solvent such as tetrahydrofuran or dioxane, at a temperature ranging between 25° and the solvent reflux, for a time from 1 to 24 hours (step (4)).

A starting compound VII can be obtained, for example, following the synthetic sequence shown in scheme 3.

12

Scheme 3



In this sequence the starting product is a compound of general formula XII, wherein R^2 represents the groups described above and P represents a suitable hydroxy-protecting group, such as a methyl or t-butyldimethylsilyl group. Said compound can be obtained easily starting from the corresponding hydroxynaphthoic acids, following preparation processes described in literature (Gray G.W. et al., J. Chem. Soc., 678 10 (1954) and Daines et al., WO 9217172).

Starting from XII, VIIa (VII with $m=0$) can be obtained removing the protecting group P (step (5)); for example when this is a methyl group, it can be removed by means of boron tribromide in an organic solvent such as methylene chloride or chloroform, at a temperature ranging between -78° and 0°C for a time from 2 to 8 hours. When P is a t-butyldimethylsilyl group, it can be removed with a base such as potassium carbonate or bicarbonate, in a solvent such as tetrahydrofuran, methanol or dioxane at a temperature ranging between 0° and 50°C , for a time from 2 to 12 hours.

A compound VIIb (VII with $R^1=\text{H}$ and $m=1$) can be obtained also starting from a compound XII. By means of a step (6), a compound XII is subjected to the action of a suitable metal hydride, such as sodium borohydride, in an solvent such as methanol, ethanol or tetrahydrofuran, in the presence of a catalytic amount of water, at a temperature ranging between 20°C and the solvent reflux, for a time from 3 to 24 hours: in this way compound XIII is obtained. This compound is subjected to a tosylation reaction (step (7)) with tosyl chloride in the presence of pyridine or triethylamine in methylene chloride, to


give a compound of formula XIV, which is treated with sodium cyanide in tetrahydrofuran or DMSO at a temperature between 25°C and the solvent reflux to yield compound XV (step (8)). The hydrolysis of XV, for example with NaOH in ethanol at a temperature ranging between 25°C and the solvent reflux, followed by the esterification with methanol in the presence of sulfuric acid, gives compound XVII (steps (9) and (10) respectively). Finally, the removal of the protecting group P, as described above, can lead to a compound VIIb.

Analogously, a compound VIIc (VII with $R^1=CH_3$ and $m=1$) can be obtained by a synthesis sequence in which the starting product is a compound XV. By treating XV with a strong base, such as sodium methoxide or potassium t-butoxide, in a suitable solvent such as tetrahydrofuran or N,N-dimethylformamide, at a temperature ranging between -30° and 25°C for a time from 2 to 24 hours (step (12)), a compound XVIII is obtained. After that, by hydrolysis of the nitrile group present in XVIII (step (13)), esterification of the carboxylic group of XIX (step (14)) and elimination of the protecting group P in XX (step (15)), as described above, compound VIIc is obtained.

Alternatively, a compound VIIb can be obtained following the synthesis shown in scheme 4.

15

Scheme 4



VIIb (VII with R¹=H and m=1)

Starting from a compound of general formula XXI, which is prepared easily following processes described in literature (Müller et al. *Helv. Chim. Acta*, 57, 790 (1974)), a compound XXII can be prepared by the modified Willgerodt's reaction. By means of such a reaction, treating a compound XXI with sulfur in the presence of an amine such as morpholine, which in its turn acts as the solvent, at the solvent reflux temperature, for a time from 6 to 24 hours, a compound of formula XXII (step (16)) is obtained after a suitable treatment with hydrochloric acid. The removal of the hydroxy-protecting methyl group, according to the process described above, leads to the preparation of a compound VIIb (step (17)).


Alternatively, a compound VIId, equivalent to VII with R¹=CH₃, R²=H, m=1 and with an alkylmethoxycarbonyl substituent at the 6- position of the 2-naphthol system can be prepared, starting from a compound XXIII, which is the known NSAID naproxen, commercially available both as the racemate and in the form of the two resolved enantiomers. Compound XXIII is subjected to demethylation with BBr₃ in a solvent such as chloroform

16

or methylene chloride at a temperature between -78°C and room temperature for 1 to 6 hours, followed by addition of methanol to the reaction, to obtain the desired compound VIId (scheme 5).


5

Scheme 5



15 Alternatively a compound IIa, i.e. a compound II with $m=0$, can be obtained following the process described in scheme 6.

Scheme 6


IIa (II with $m=0$)

30 Starting from a compound XXIV, easily available following chemical processes described in literature (Tolbert et al. J. Am. Chem. Soc., 112, 8163 (1990)), a

compound XXV can be prepared by condensation with VIII (step (18)) following the method described for the preparation of the compounds of general formula IX. The hydrolysis of XXV (step (19)), according to the process described for the preparation of XIX, leads to the compounds of general formula IIa.

A compound of general formula Xa, i.e. of general formula X with $R^1=H$, $R^4=H$ and $m=1$, can be obtained, for example, following the synthetic sequence shown in scheme 7.

Scheme 7



A compound XXVII can be obtained starting from a compound of general formula XXVI, commercial or easily available starting from XXIV, by catalytic hydrogenation with a suitable catalyst, such as Pd-C or Pd(OH)₂-C in a suitable solvent such as ethanol or methanol, in the presence of an acid such as acetic acid or hydrochloric

acid, under hydrogen pressures ranging between atmosphere pressure and 2 kg/cm^2 , at a temperature from 25° to 50°C , for a time from 3 to 24 hours (step (20)).


5 A compound XXVIII can be obtained by acetylation of XXVII for example with acetic anhydride in the presence of a base such as pyridine or triethylamine in a solvent such as chloroform or methylene chloride, at a temperature between -78° and 25°C for a time from 1 to 6 hours (step (21)).

10 A compound Xa can be obtained starting from XXVIII by deprotecting the methylated hydroxy group (step (22)) according to the process described above. Analogously, a compound Xb, i.e. a compound of general formula X with $R^1=\text{H}$, $R^4=\text{CH}_3$ and $m=1$, can be obtained by methylation of
15 a compound XXVIII (step (23)) with methyl iodide or methyl sulfate in the presence of a base such as lithium diisopropylamide, lithium amide or potassium t-butoxide in a suitable solvent such as tetrahydrofuran, benzene or toluene at a temperature ranging between -78° and
20 25°C for 2-6 hours. The subsequent deprotection of the methylated hydroxy group leads to a compound Xb (step (24)).

25 A compound Xc, i.e. a compound of general formula X with $R^1=\text{CH}_3$ and $m=1$, can be obtained for example following the synthetic sequence shown in scheme 8.

19

Scheme 8



The reduction of a compound of formula XXX (easily available starting from XXI), for example, with a suitable hydride such as sodium borohydride in a solvent such as methanol, ethanol or tetrahydrofuran at a temperature between 20° and the solvent reflux for a time from 3 to 24 hours, gives a compound XXXI (step (25)).

A compound XXXII can be obtained, for example, by treatment of a compound XXXI (step (26)) with p-toluenesulfonic acid chloride in the presence of a base such as pyridine or triethylamine which can in their turn act as solvents, the presence of other solvents such as chloroform or methylene chloride being optional, at a temperature ranging between 20° and 40°C for a time from 3 to 18 hours.

A compound of formula XXXIII can be prepared for example starting from a compound XXXII (step (27)) by


replacing the tosylate group with an azido group, using sodium azide as the reactive, in a suitable organic solvent such as N,N-dimethylformamide or tetrahydrofuran at a temperature ranging between 20°C and the solvent
5 reflux for a time from 6 to 24 hours; and following catalytic hydrogenation of the intermediate azide with a suitable catalyst such as Pd-C or in a solvent such as ethanol, methanol or ethyl acetate, under hydrogen pressures ranging between atmosphere pressure and 2
10 kg/cm², at a temperature between 0° and 50°C, for a time from 6 to 24 hours.

Finally a compound Xc can be obtained by successive steps of acetylation, optional methylation of the amido group and deprotection of the methylated hydroxyl, as
15 described above for the compounds Xa and Xb.

Analogously, a compound Xd, i.e. of general formula X with m=0, can be prepared starting from XXXV, easily available according to processes described in literature (Airan et al., J. Am. Chem. Soc., 28, 339 (1927)), by
20 the process shown in scheme 9, in which a synthesis sequence similar to that described for the preparation of the other compounds Xa-c is used.


21

Scheme 9



A starting compound III can be prepared following the synthesis sequence shown in scheme 10.

Scheme 10



Starting from a compound XXXVIII, commercial or easily available according to similar chemical processes, wherein m and p have the values described above, R⁵ represents the groups described above, except hydrogen and X can be a halogen atom, a compound XXXIX can be prepared by treatment with NaCN in a suitable solvent such as dimethylsulfoxide, water or ethanol, at a temperature ranging between 25°C and the solvent reflux for 1-8 hours (step 33). By reduction of the cyano group in XXXIX, following the same process as described in step (20), a compound IIIa can be prepared, which is equivalent to a compound of general formula III with R⁴=H and D=COOR⁵.

Starting from a compound IIIa, a compound IIIb can be obtained, i.e. of general formula III with R⁴=H and D=5-tetrazolyl, by a process which comprises the steps (36)-(40). First, the amino group in IIIa is protected with a benzyloxycarbonyl or t-butoxycarbonyl group, according to processes widely described in literature, to yield compound XL, wherein Q represents one of the amino-protecting groups mentioned above. The basic hydrolysis of XL leads to the preparation of XLI, starting from which a compound XLII can be prepared by means, for example, of a reaction with ethyl chloroformate in the presence of a base such as triethylamine or pyridine in a solvent such as tetrahydrofuran or ethyl ether and following treatment with ammonia, at a temperature between 0° and 25°C, for a time from 30 min to 3 hours. The dehydration of a compound XLII, for example with phosphorous oxychloride, in a solvent such as N,N-dimethylformamide at a

temperature ranging between 0° and 50°C for 3-24 hours, gives a compound XLIII. The treatment of a compound XLIII with sodium azide in a suitable solvent such as N,N-dimethylformamide at a temperature ranging between
5 25°C and the solvent reflux and the subsequent elimination of the protective group present in the amino group, according to conventional techniques, yields a compound IIIf. Alternatively, when a compound XLIV is commercial or easily available by similar synthetic
10 methods, a compound IIIf can be obtained starting from XLIV by reaction with sodium azide (step (42)) and subsequent reduction of the azide (step 43), as described above in step (27).

Starting from a compound of formula IIIf or of
15 formula IIIf, a compound IIIfc or IIIfd can be obtained respectively by methylation of the amine according to, for example, a process comprising first the formylation of the amino group, with acetic anhydride and formaldehyde mixtures in a suitable solvent, such as
20 tetrahydrofuran or ethyl ether, at a temperature between 0° and 25°C for 3-24 h, followed by reduction of the formyl group with the BH₃-tetrahydrofuran complex in a solvent such as tetrahydrofuran or ethyl ether at a temperature ranging between -78° and 0°C for 6-24 hours.

25 A starting compound of formula VI can be prepared for example following the process shown in scheme 11.


Starting from a compound XLVI, wherein m and p have the meanings described above and R⁵ also represents the groups described above, except hydrogen, and Y
30 represents a chlorine or bromine atom, commercially available in some instances or easily prepared by

similar synthetic methods, VIa, i.e. the compound of general formula VI with $D=COOR^5$, (steps (44) and (45)), can be obtained following the processes described above for steps (33) and (13).


5

Scheme 11


Y=Cl o Br




10



XLVIII

VIa (VI with $D=COOR^5$)

15

20. Vlb (VI with $D=5\text{-tetrazolyl}$)

Starting from XLVI, wherein Y represents a hydroxyl group, a compound XLVIII can be prepared following a process similar to that described in scheme 10. Finally, a compound Vlb, i.e. of general formula VI with $D=5\text{-tetrazolyl}$, can be prepared by a process which comprises the tosylation of XLVIII, the substitution of the tosylate by a nitrile group and the final hydrolysis according to the above described methods.


25

Alternatively, when in VI $m=0$, $D=COOR^5$ and the two substituents of the benzene ring are in para position, compound of formula VIc can be prepared following the

30

synthesis shown in scheme 12.

Scheme 12



10 VIc (VI with m=0, D=COOR⁵
and para orientation
of the substituents)

A compound L can be prepared starting from XLIX, for example, by a formylation reaction using a suitable reactive such as hexamethylenetetramine or N,N-dimethylformamide in the presence of trifluoroacetic acid or phosphorous oxychloride at a temperature ranging between 25° and 100°C, for a time from 2 to 24 hours. The subsequent oxidation with a suitable oxidizing agent such as Jones's reagent at a temperature ranging between 0° and 25°C for a time of 2 to 18 hours, leads to the desired compound VIc.

25 The compounds of the present invention show a marked antagonistic activity of leukotrienes effects and they have therefore anti-inflammatory and anti-allergic properties which make them useful in the treatment of diseases wherein those mediators are involved.

30 Said compounds can be therefore used in human therapy, for the prevention and treatment of allergic rhinitis, bronchial asthma, hypersensitivity reactions such as allergic conjunctivitis, various inflammatory conditions such as rheumatoid arthritis, osteoarthritis,

tendinitis, bursitis, psoriasis and related inflammations.

The compound of the present invention may also be used in the treatment of diseases of the cardiovascular system, such as cardiac ischemia, myocardic infarct, coronary spasm, cardiac anaphylaxis, cerebral oedema and endotoxic shock.

For the intended therapeutic uses, the compounds of the invention are formulated in suitable pharmaceutical compositions, using conventional techniques and methods, as disclosed in Remington's Pharmaceutical Science Handbook, Mack Pub. Co., N.Y. U.S.A. Examples of said formulations include capsules, tablets, syrups and the like, containing from 1 to 1000 mg of active principle per unit dose.

EXAMPLES

The following examples illustrate the preparation and the pharmacological activity of the compounds of the present invention.

Example 1: N-[4-(1H-5-tetrazolyl)phenylmethyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide
1A Methyl 2-(6-hydroxy-2-naphthyl)propionate

Boron tribromide (70 ml, 737 mmol) was added at -78°C to a solution of 2-(6-methoxy-2-naphthyl)propionic acid (10 g, 43.4 mmol) in dry methylene chloride (130 ml). The reaction mixture was stirred at room temperature for 5 h, then 136 ml of methanol were added and stirred for 18 hours. After this time, the mixture was evaporated to dryness, water (250 ml) was added and extracted with ethyl ether (4x100 ml). The combined ether phases were dried and the solvent was evaporated

off, to obtain a crude which was purified by flash chromatography through a silica gel column. Eluting with petroleum ether:ethyl ether, 9:1, 6.9 g of the title compound were obtained as a white solid with melting point 184-186°C (70% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.56 (d, 3H); 3.70 (s, 3H); 3.87 (q, 1H); 5.75 (s, 1H); 7.08 (m, 2H); 7.38 (dd, 1H); 7.60 (d, 1H); 7.66 (m, 2H).

1B Methyl 2-[6-(2-quinolinylmethoxy)-2-naphthyl]propionate

Sodium methoxide (6.0 ml, 31.2 mmol) was added to a solution of methyl 2-(6-hydroxy-2-naphthyl)propionate (7.2 g, 31.2 mmol) in DMF (200 ml) and stirred at room temperature for 5 min. After that 2-chloromethylquinoline (5.5 g, 31.2 mmol) was added thereto and the reaction mixture was stirred at room temperature for 18 h, then evaporated to dryness, the residue was dissolved in ethyl acetate (250 ml), washed with 5% NaHCO₃ (3x25 ml), dried and the solvent was evaporated off, to obtain a crude which was purified by crystallization with methanol. 9.3 g of the title compound were obtained as a white solid with melting point 98-99°C (80% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.42 (d, 3H); 3.53 (s, 3H); 3.87 (q, 1H); 5.44 (s, 2H); 7.30 (m, 2H); 7.39 (d, 1H); 7.56 (t, 1H); 7.67 (m, 5H); 7.93 (d, 1H); 7.99 (d, 1H); 8.37 (d, 1H).

1C 2-[6-(2-Quinolinylmethoxy)-2-naphthyl]propionic acid

1M lithium hydroxide (31.7 ml) was added to a solution of methyl 2-[6-(2-quinolinylmethoxy)-2-naphthyl]propionate (5 g, 13.2 mmol) in THF (40 ml) and

stirred at room temperature for 48 h. After that THF was evaporated off, pH was adjusted to 4-5 with 1M HCl and the mixture was extracted with ethyl acetate, to obtain 4.7 g of the title compound as a white solid with melting point 192-194°C (99% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.39 (d, 3H); 3.76 (q, 1H); 5.44 (s, 2H); 7.29 (dd, 1H); 7.35 (dd, 1H); 7.40 (d, 1H); 7.58 (t, 1H); 7.69 (m, 3H); 7.77 (dt, 1H); 7.80 (d, 1H); 7.95 (d, 1H); 7.98 (d, 1H); 8.38 (d, 1H).

10 1D 4-(1H-5-Tetrazolyl)azidomethylbenzene

Sodium azide (15 g, 230 mmol) and ammonium chloride (12.3 g, 230 mmol) were added to a solution of 4-bromomethylbenzonitrile (5 g, 25.5 mmol) in DMF (75 ml). The reaction mixture was stirred at 110°C for 16 h, then poured onto 200 ml of 1M HCl and extracted with ethyl acetate (4x75 ml). The organic phase was dried and solvents were removed, to obtain a oil which was diluted with ethyl ether and petroleum ether, to give a precipitate which was filtered and washed with petroleum ether. In this way 5 g of the title compound were obtained (97% yield).

¹H N.M.R. (300 MHz, CD₃OD) δ ppm: 4.52 (s, 2H); 7.59 (d, 2H); 8.07 (d, 2H).

15 1E 4-(1H-5-Tetrazolyl)benzylamine hydrochloride

25 1.4 g of 10% palladium on charcoal was added to a solution of 4-(1H-5-tetrazolyl)azidomethyl-benzene (4.3 g, 21.5 mmol) in methanol (400 ml) and concentrated HCl (14 ml) and stirred at room temperature for 4 days, under hydrogen atmosphere. After that the reaction mixture was filtered and the filtrate was evaporated to dryness to obtain a crude which was redissolved in hot

30

methanol, crystallizing in this way 3.7 g of the title compound as a yellowish solid with melting point $>360^{\circ}\text{C}$ (80% yield).

5 ^1H N.M.R. (300 MHz, DMSO) δ ppm: 4.16 (d, 2H); 7.75 (d, 2H); 8.18 (d, 2H); 8.56 (s, 3H).

1F N-[4-(1H-5-Tetrazolyl)phenylmethyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide

4-(1H-5-tetrazolyl)benzylamine hydrochloride (0.212 g, 1.0 mmol), N,N-dimethylaminopyridine (0.373 g, 3.05 mmol) and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.202 g, 1.05 mmol) were added in succession to a solution of 2-[6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid (0.376 g, 1.05 mmol) in dry methylene chloride (50 ml).
15 The reaction mixture was stirred at room temperature for 24 h, thereafter was poured onto water (30 ml) and pH was adjusted to 5 with 1M HCl, the phases were separated and the organic one was extracted with ethyl acetate (4x50 ml). The combined organic extracts were dried and
20 the solvent was evaporated off, to obtain a crude which was purified by crystallization in methanol-chloroform-acetic acid mixtures, thereby obtaining 0.351 g of the title compound as a white solid with melting point $214-215^{\circ}\text{C}$ (65% yield).

25 ^1H N.M.R. (300 MHz, DMSO) δ ppm: 1.45 (d, 3H); 3.83 (q, 1H); 4.33 (d, 2H); 5.49 (s, 2H); 7.30 (dd, 1H); 7.35 (d, 2H); 7.44 (d, 1H); 7.47 (dd, 1H); 7.63 (dt, 1H); 7.72 (s, 1H); 7.78 (m, 4H); 7.92 (d, 2H); 8.01 (d, 1H); 8.05 (d, 1H); 8.44 (d, 1H); 8.61 (t, 1H).

30 Example 2: N-[3-(1H-5-tetrazolyl)phenylmethyl]-2-

[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide

Following the process described in example 1 (point F), starting from 2-[6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid and 3-(1H-5-tetrazolyl)benzylamine hydrochloride, the title compound was prepared as a white solid with melting point 114-115°C (70% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.48 (d, 3H); 3.86 (q, 1H); 4.38 (m, 2H); 5.52 (s, 2H); 7.33 (dd, 1H); 7.47 (m, 4H); 7.66 (dt, 1H); 7.80 (m, 6H); 7.98 (s, 1H); 8.04 (d, 1H); 8.09 (d, 1H); 8.47 (d, 1H); 8.66 (t, 1H).

Example 3: N-[2-(1H-5-tetrazolyl)phenylmethyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide

Following the process described in example 1 (point F) starting from 2-[6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid and 2-(1H-5-tetrazolyl)benzylamine hydrochloride, the title compound was prepared as a yellowish solid with melting point 189-190°C (57% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.40 (d, 3H); 3.80 (q, 1H); 4.52 (d, 2H); 5.49 (s, 2H); 7.35 (m, 6H); 7.63 (dt, 1H); 7.76 (m, 6H); 8.01 (dd, 1H); 8.05 (d, 1H); 8.43 (d, 1H); 8.49 (t, 1H).

Example 4: N-(4-cyanomethylphenyl)-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide

Following the process described in example 1 (point F), starting from 2-[6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid and 4-cyanomethylaniline, the title compound was prepared as a white solid (recrystallized from methanol) with melting point 177-178°C (77% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.45 (d, 3H); 3.90 (m, 3H); 5.43 (s, 2H); 7.20 (d, 2H); 7.27 (dd, 1H); 7.39 (d, 1H); 7.46 (d, 1H); 7.56 (m, 3H); 7.69 (t, 2H); 7.48 (m, 2H); 7.81 (d, 1H); 7.94 (d, 1H); 8.00 (d, 1H); 8.36 (d, 1H); 10.13 (s, 1H).

Example 5: N-[4-(1H-5-tetrazolyl)methylphenyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide

Following the process described in example 1 (point D), starting from N-(4-cyanomethylphenyl)-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide, the title compound was prepared as a white solid (recrystallized from methanol-ethyl acetate) with melting point 179.1-180.1°C (88% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.47 (d, 3H); 3.93 (q, 1H); 4.20 (s, 2H); 5.47 (s, 2H); 7.16 (d, 2H); 7.30 (dd, 1H); 7.42 (d, 1H); 7.49 (dd, 1H); 7.55 (d, 2H); 7.60 (dt, 1H); 7.75 (m, 4H); 7.83 (d, 1H); 7.97 (d, 1H); 8.03 (d, 1H); 8.40 (d, 1H); 10.11 (s, 1H).

Example 6: N-[4-(1H-5-tetrazolyl)methylphenyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide (sodium salt)

1M sodium hydroxide (0.31 ml) was added to a solution of N-[4-(1H-5-tetrazolyl)methylphenyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide (0.161 g, 0.31 mmol) in methanol (5 ml) and stirred at room temperature for 30 min. After that the reaction mixture was evaporated to dryness and the residue was recrystallized from ethanol-ethyl ether, to obtain 0.143 g of the title compound as a white solid with melting point 278-279°C (85% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.47 (d, 3H); 3.93 (m,

3H); 5.47 (s, 2H); 7.16 (d, 2H); 7.30 (dd, 1H); 7.42 (d, 1H); 7.49 (dd, 1H); 7.55 (d, 2H); 7.60 (dt, 1H); 7.75 (m, 4H); 7.83 (d, 1H); 7.97 (d, 1H); 8.03 (d, 1H); 8.40 (d, 1H); 10.11 (s, 1H).

5 Example 7: methyl 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidolbenzoate

Following the process described in example 1 (point F), starting from 2-[6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid and methyl 4-amino-
10 benzoate, the title compound was prepared as a white solid with melting point 182.0-182.3°C (72% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.58 (dd, 3H); 3.81 (m, 4H); 5.43 (s, 2H); 7.18 (s, 1H); 7.24 (m, 1H); 7.35 (dd, 1H); 7.50 (m, 3H); 7.75 (m, 8H); 8.10 (d, 1H); 8.17
15 (d, 1H); 10.41 (s, 1H).

Example 8: 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidolbenzoic acid

Following the process described in example 1 (point C), starting from methyl 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidolbenzoate, the title
20 compound was prepared as a white solid (recrystallized from ethanol) which decomposes at 264-265°C (74% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.49 (d, 3H); 4.00 (q, 1H); 5.48 (s, 2H); 7.32 (dd, 1H); 7.42 (d, 1H); 7.58 (d, 1H); 7.75 (m, 10H); 7.99 (d, 1H); 8.04 (d, 1H); 8.42 (d, 25
1H); 10.41 (s, 1H).

Example 9: methyl 4-[4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidolphenyl]butanoate

9A Methyl 4-(4-aminophenyl)butanoate

30 8.4 ml of concentrated H₂SO₄ was added to a solution of 4-(4-aminophenyl)butanoic acid (2.0 g, 11.6

mmo) in methanol (84 ml) and refluxed for 2 h. After that the mixture was left to cool at room temperature, added with Na_2CO_3 to basic pH and extracted with ethyl acetate. The organic phase was dried and the solvent was
5 evaporated off, to obtain 1.7 g of the title compound as a colourless oil (82% yield).

^1H N.M.R. (300 MHz, CD_3OD) δ ppm: 1.83 (q, 2H); 2.28 (t, 2H); 2.49 (t, 2H); 3.62 (s, 3H); 6.66 (d, 2H); 6.92 (d, 2H).

10 9B Methyl 4-[4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidolphenyl]butanoate

Following the process described in example 1 (point F), starting from 2-[6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid and methyl 4-(4-amino-
15 phenyl)butanoate, the title compound was prepared as a white solid with melting point 145.7-149.0°C (87% yield).

^1H N.M.R. (300 MHz, CDCl_3) δ ppm: 1.61 (d, 3H); 1.82 (q, 2H); 2.27 (t, 2H); 2.56 (t, 2H); 3.62 (s, 3H); 3.81 (m, 1H); 5.42 (s, 2H); 7.02 (d, 2H); 7.28 (dd, 1H); 7.37 (m, 3H); 7.70 (m, 7H); 7.81 (d, 1H); 8.12 (d, 1H); 8.19 (d, 1H).
20

Example 10: 4-[4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidolphenyl]butanoic acid

25 Following the process described in example 1 (point C), starting from methyl 4-[4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamido]phenyl]butanoate, the title compound was prepared as a white solid with melting point 176.0-176.3°C (68% yield).

30 ^1H N.M.R. (300 MHz, DMSO) δ ppm: 1.42 (d, 3H); 1.68 (q, 2H); 2.12 (t, 2H); 2.45 (t, 2H); 3.97 (q, 1H); 5.55 (s,

2H); 7.03 (d, 2H); 7.26 (dd, 1H); 7.38 (d, 1H); 7.48 (m, 3H); 7.57 (t, 1H); 7.72 (m, 5H); 7.95 (d, 1H); 8.00 (d, 1H); 8.38 (d, 1H).

Example 11: methyl 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidomethyl]benzoate

11A Methyl 4-cyanobenzoate

Following the process described in example 9 (point A), starting from 4-cyanobenzoic acid, the title compound was prepared as a yellowish oil (93% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 4.11 (s, 3H); 7.89 (d, 2H); 8.28 (d, 2H).

11B Methyl 4-aminomethylbenzoate

Following the process described in example 1 (point E), starting from methyl 4-cyanobenzoate and reacting for 4 h, the title compound was prepared as a semi-solid oil (87% yield).

¹H N.M.R. (300 MHz, CD₃OD) δ ppm: 3.91 (s, 3H); 4.21 (s, 2H); 7.59 (d, 2H); 8.07 (d, 2H).

11C Methyl 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidomethyl]benzoate

Following the process described in example 1 (point F), starting from 2-(6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid and methyl 4-aminomethylbenzoate, the title compound was prepared as a white solid with melting point 149.0-150.8°C (68% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.09 (d, 3H); 3.79 (s, 3H); 3.85 (q, 1H); 4.32 (s, 2H); 5.61 (s, 2H); 7.08 (m, 4H); 7.30 (m, 2H); 7.47 (m, 2H); 7.62 (m, 3H); 7.74 (t, 1H); 7.82 (d, 1H); 7.89 (d, 1H); 8.45 (d, 1H); 8.52 (d, 1H).

Example 12: 4-[2-[6-(2-Quinolinylmethoxy)-2-naphthyl]-

propanamidomethylbenzoic acid

Following the process described in example 1 (point C), starting from methyl 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidomethyl]benzoate, the title compound was prepared as a white solid with melting point 203.9-205.8°C (68% yield).

¹H N.M.R. (300 MHz, CD₃OD) δ ppm: 1.55 (d, 3H); 3.85 (q, 1H); 4.40 (q, 1H); 5.66 (s, 2H); 7.22 (d, 2H); 7.36 (dd, 1H); 7.41 (d, 1H); 7.47 (dd, 1H); 7.80 (m, 6H); 8.01 (m, 2H); 8.16 (d, 1H); 8.24 (d, 1H); 8.81 (d, 1H).

Example 13: methyl 3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidomethyl]benzoate13A Methyl 3-cyanobenzoate

Following the process described in example 9 (point A), starting from 3-cyanobenzoic acid, the title compound was prepared as a yellowish oil (80% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 3.93 (s, 3H); 7.56 (t, 1H); 7.81 (dd, 1H); 8.23 (dd, 1H); 8.29 (d, 1H).

13B Methyl 3-aminomethylbenzoate

Following the process described in example 1 (point E), starting from methyl 3-cyanobenzoate and reacting for 4 h, the title compound was prepared as a semi-solid oil (92% yield).

¹H N.M.R. (300 MHz, CD₃OD) δ ppm: 3.88 (s, 3H); 3.91 (s, 2H); 7.49 (t, 1H); 7.51 (dd, 1H); 7.91 (dd, 1H); 7.99 (d, 1H).

13C Methyl 3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidomethyl]benzoate

Following the process described in example 1 (point F), starting from 2-[6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid and methyl 3-aminome-

thylbenzoate, the title compound was prepared as a white solid with melting point 135.4-136.8°C (63% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.56 (d, 3H); 3.69 (q, 1H); 3.79 (s, 3H); 4.36 (d, 2H); 5.43 (s, 2H); 7.16 (d, 1H); 7.27 (m, 4H); 7.50 (t, 1H); 7.70 (m, 8H); 8.05 (d, 1H); 8.13 (d, 1H).

Example 14: 3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]-propanamidomethyl]benzoic acid

Following the process described in example 1 (point C), starting from methyl 3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidomethyl]benzoate, the title compound was prepared as a white solid with melting point 203.5-204.7°C (77% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.43 (d, 3H); 3.80 (q, 1H); 4.27 (q, 2H); 7.25 (m, 3H); 7.42 (m, 2H); 7.62 (m, 1H); 7.75 (m, 5H); 8.00 (d, 1H); 8.05 (d, 1H); 8.32 (s, 1H); 8.42 (d, 1H); 8.56 (t, 1H).

Example 15: methyl 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidolphenylacetate

15A Methyl 4-aminophenylacetate

Following the process described in example 9 (point A), starting from 4-aminophenylacetic acid, the title compound was prepared as a yellowish oil (73% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 3.50 (s, 2H); 3.65 (s, 3H); 3.68 (s, 2H); 6.59 (d, 2H); 7.04 (d, 2H).

15B Methyl 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]-propanamidolphenylacetate

Following the process described in example 1 (point F), starting from 2-(6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid and methyl 4-aminophenylacetate, the title compound was prepared as a

white solid with melting point 154.0-156.0°C (80% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.64 (d, 3H); 3.53 (s, 2H); 3.64 (s, 3H); 3.83 (q, 1H); 5.50 (s, 2H); 7.03 (m, 1H); 7.15 (d, 2H); 7.31 (m, 4H); 7.56 (t, 1H); 7.72 (m, 5H); 7.83 (d, 1H); 8.11 (d, 1H); 8.19 (d, 1H).

Example 16: 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]-propanamidolphenylacetic acid

Following the process described in example 1 (point C), starting from methyl 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidolphenylacetate, the title compound was prepared as a white solid with melting point 141.0-143.2°C (77% yield).

¹H N.M.R. (300 MHz, CDCl₃-CD₃OD) δ ppm: 1.69 (d, 3H); 3.63 (s, 2H); 4.04 (q, 1H); 5.60 (s, 2H); 7.28 (d, 2H); 7.39 (d, 2H); 7.59 (d, 3H); 7.72-7.80 (m, 2H); 7.86-7.96 (m, 4H); 8.05 (d, 1H); 8.22 (d, 1H); 8.53 (d, 1H).

Example 17: methyl 3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidolphenylacetate

17A Methyl 3-aminophenylacetate

Following the process described in example 9 (point A), starting from 3-aminophenylacetic acid, the title compound was prepared as a yellowish oil (81% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 3.52 (s, 2H); 3.67 (s, 3H); 3.68 (s, 2H); 6.55 (m, 2H); 6.65 (d, 1H); 7.09 (t, 1H).

17B Methyl 3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]-propanamidolphenylacetate

Following the process described in example 1 (point F), starting from 2-(6-(2-quinolinylmethoxy)-2-naphthyl]propionic acid and methyl 3-amino-

phenylacetate, the title compound was prepared as a white solid with melting point 141.8-142.8°C (75% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.65 (d, 3H); 3.54 (s, 2H); 3.64 (s, 3H); 3.82 (q, 1H); 5.51 (s, 2H); 6.96 (d, 1H); 7.03 (s, 1H); 7.35 (m, 6H); 7.56 (t, 1H); 7.75 (m, 6H); 8.12 (d, 1H).

Example 18: 3-[2-[6-(2-Quinolinylmethoxy)-2-naphthyl]propanamidolphenylacetic acid

Following the process described in example 1 (point C), starting from methyl 3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidolphenylacetate, the title compound was prepared as a white solid with melting point 200.9-202.9°C (61% yield).

¹H N.M.R. (300 MHz, CDCl₃-CD₃OD) δ ppm: 1.59 (d, 3H); 3.55 (s, 2H); 3.97 (q, 1H); 5.79 (s, 2H); 6.99 (d, 1H); 7.22 (t, 1H); 7.45 (m, 6H); 7.77 (d, 1H); 7.83 (m, 2H); 7.90 (m, 1H); 8.17 (m, 2H); 8.31 (dd, 1H); 8.37 (d, 1H); 9.1 (m, 1H).

Example 19: methyl 4-[4-[2-[6-[(7-chloro-2-quinoliny)lmethoxy]-2-naphthyl]propanamidolphenyl]butanoate
19A 2-Bromomethyl-7-chloroquinoline

N-bromosuccinimide (3.3 g, 18.5 mmol) and some crystals of 2,2'-azobis(2-methylpropionitrile) (AIBN) were added to a solution of 7-chloro-2-methylquinoline (3 g, 16.9 mmol) in dry carbon tetrachloride (100 ml). The reaction mixture was refluxed for 4 h, then cooled at room temperature. The precipitate was filtered off and the filtrate was washed with a NaCl saturated solution (3x20 ml), dried and the solvent was evaporated off to obtain a crude which was

purified by flash chromatography through a silica gel column. Eluting with petroleum ether:chloroform, 3:2, 2.3 g of the title compound were obtained as a white solid with melting point 110-111°C (52% yield).

5 ¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 4.65 (s, 2H); 7.47 (dd, 1H); 7.53 (d, 1H); 7.71 (d, 1H); 8.03 (d, 1H); 8.11 (d, 1H).

19B Methyl 2-[6-(7-chloro-2-quinolinyl)methoxy-2-naphthyl]propionate

10 Following the process described in example 1 (point B), starting from methyl 2-(6-hydroxy-2-naphthyl)propionate and 2-bromomethyl-7-chloroquinoline, the title compound was prepared as a white solid with melting point 98-99°C (78% yield)

15 ¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.56 (d, 3H); 3.65 (s, 3H); 3.82 (q, 1H); 5.47 (s, 2H); 7.19 (d, 1H); 7.25 (s, 1H); 7.28 (dd, 1H); 7.37 (dd, 1H); 7.50 (dd, 1H); 7.70 (m, 4H); 8.10 (d, 1H); 8.14 (d, 1H).

19C 2-[6-(7-Chloro-2-quinolinyl)methoxy-2-naphthyl]propanoic acid

20 Following the process described in example 1 (point C), starting from methyl 2-[6-(7-chloro-2-quinolinyl)methoxy-2-naphthyl]propionate, the title compound was prepared (90% yield).

25 ¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.48 (d, 3H); 3.70 (q, 1H); 5.49 (d, 2H); 7.25 (m, 2H); 7.49 (dd, 1H); 7.58 (dd, 1H); 7.63 (d, 1H); 7.77 (m, 3H); 7.93 (d, 1H); 8.05 (d, 1H); 8.37 (d, 1H).

19D Methyl 4-[4-[2-[6-[(7-chloro-2-quinolinyl)methoxy]-2-naphthyl]propanamidolphenyl]butanoate

30 Following the process described in example 1 (point

F), starting from 2-[6-(7-chloro-2-quinolinyloxy)methoxy-2-naphthyl]propanoic acid and methyl 4-(4-aminophenyl)butanoate, the title compound was prepared as a white solid with melting point 131.0-133.0°C (60% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.58 (d, 3H); 1.80 (q, 2H); 2.20 (t, 2H); 2.48 (t, 2H); 3.56 (s, 3H); 3.75 (q, 1H); 5.41 (s, 2H); 7.06 (m, 3H); 7.23 (dd, 1H); 7.30 (m, 3H); 7.41 (dd, 1H); 7.50 (dd, 1H); 7.73 (m, 4H); 8.10 (d, 1H); 8.16 (d, 1H).

Example 20: 4-[4-[2-[6-[(7-chloro-2-quinolinyloxy)methoxy]-2-naphthyl]propanamido]phenyl]butanoic acid

Following the process described in example 1 (point C), starting from methyl 4-[4-[2-[6-[(7-chloro-2-quinolinyloxy)methoxy]-2-naphthyl]propanamido]phenyl]butanoate, the title compound was prepared as a white solid with melting point 176.4-177.8°C (69% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.42 (d, 3H); 1.69 (q, 2H); 2.11 (t, 2H); 3.89 (q, 1H); 5.43 (s, 2H); 7.03 (d, 2H); 7.26 (dd, 1H); 7.37 (d, 1H); 7.45 (m, 3H); 7.61 (dd, 1H); 7.71 (m, 3H); 7.81 (d, 1H); 8.01 (d, 1H); 8.05 (s, 1H); 8.42 (d, 1H); 9.99 (s, 1H).

Example 21: N-[4-(1H-5-tetrazolyl)phenylmethyl]-6-(2-quinolinyloxy)-2-naphthaleneacetamide

21A 2-Acetyl-6-methoxynaphthalene

2-methoxynaphthalene (10 g, 63.3 mmol) followed by acetyl chloride (5.8 ml, 79.8 mmol) were added drop by drop to a solution of aluminium trichloride (10.9 g, 81.7 mmol) in nitrobenzene (50 ml) cooled at 0°C and under inert atmosphere. The reaction mixture was stirred at 0°C for 2 h and at room temperature for 18 h, then it

was cooled to 0°C, poured onto ice (50 ml), added with concentrated HCl (20 ml) and chloroform (25 ml). The two phases were separated and the organic one was washed with water (3x10 ml), dried and the solvent was evaporated off, to obtain a crude which was purified by crystallization in methanol, thereby obtaining 7.9 g of the title compound as a white solid with melting point 107-109°C (63% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 2.69 (s, 3H); 3.93 (s, 3H); 7.17 (m, 2H); 7.75 (d, 1H); 7.83 (d, 1H); 7.99 (dd, 1H); 8.37 (s, 1H).

21B Methyl 6-methoxy-2-naphthaleneacetate

A mixture consisting of 2-acetyl-6-methoxynaphthalene (3 g, 15.0 mmol), sulfur (0.72 g, 22.5 mmol) and morpholine (2 ml) was refluxed for 18 h, then acetic acid (11 ml) and concentrated HCl (18 ml) were added and reflux was continued for a further 24 h. After that the mixture was evaporated to dryness, added with methanol (60 ml) and concentrated H₂SO₄ (10 ml) and refluxed for 18 h. Finally the mixture was evaporated to dryness, added with ethyl acetate, washed with a 5% NaHCO₃ saturated solution until the washing were neutral, dried and the solvent was evaporated off, to obtain a crude, which was purified by flash chromatography through a silica gel column. Eluting with petroleum ether:chloroform, 4:1, 2.5 g of the title compound were obtained as a white solid with melting point 75.7-76.5°C (72% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 3.90 (s, 3H); 3.96 (s, 2H); 4.11 (s, 3H); 7.34 (m, 2H); 7.57 (dd, 1H); 7.85 (s, 1H); 7.88 (d, 1H); 7.90 (d, 1H).

21C Methyl 6-hydroxy-2-naphthaleneacetate

Following the process described in example 1 (point A), starting from methyl 6-methoxy-2-naphthaleneacetate, the title compound was prepared as a yellowish solid with melting point 84.8-85.8°C (72% yield)

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 3.74 (s, 3H); 3.77 (s, 2H); 6.16 (m, 1H); 7.03 (m, 2H); 7.32 (dd, 1H); 7.56 (m, 3H).

21D Methyl 6-(2-quinolinylmethoxy)-2-naphthaleneacetate

Following the process described in example 1 (point B), starting from methyl 6-hydroxy-2-naphthaleneacetate, the title compound was prepared as a white solid with melting point 106.3-107.9°C (82% yield)

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 3.72 (s, 3H); 3.78 (s, 2H); 5.64 (s, 2H); 7.26 (d, 1H); 7.33 (dd, 1H); 7.38 (dd, 1H); 7.59 (t, 1H); 7.74 (m, 5H); 7.86 (d, 3H); 8.15 (d, 1H); 8.22 (d, 1H).

21E 6-(2-Quinolinylmethoxy)-2-naphthaleneacetic acid

Following the process described in example 1 (point C), starting from methyl 6-(2-quinolinylmethoxy)-2-naphthaleneacetate, the title compound was prepared (84% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 3.75 (s, 2H); 5.60 (s, 2H); 7.20 (d, 1H); 7.31 (dd, 1H); 7.35 (dd, 1H); 7.60 (t, 1H); 7.68-7.83 (m, 6H); 8.20 (d, 1H); 8.25 (d, 1H).

21F N-[4-(1H-5-Tetrazolyl)phenylmethyl]-6-(2-quinolinylmethoxy)-2-naphthaleneacetamide

Following the process described in example 1 (point F), starting from 6-(2-quinolinylmethoxy)-2-naphthaleneacetic acid and 4-(1H-5-tetrazolyl)benzylamine hydrochloride, the title compound was

prepared as a white solid with melting point 256-257°C (58% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 3.66 (s, 2H); 4.39 (d, 2H); 5.31 (s, 2H); 7.40 (m, 5H); 7.67 (t, 1H); 7.82 (m, 6H); 7.99 (m, 3H); 8.47 (d, 1H); 8.73 (t, 1H).

Example 22: methyl 4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]carboxamidolphenyl]butanoate

22A 2-Cyano-6-hydroxynaphthalene

Following the process described in example 1 (point A), starting from 2-cyano-6-methoxynaphthalene, the title compound was prepared (91% yield).

¹H N.M.R. (300 MHz, CD₃OD) δ ppm: 7.17 (m, 2H); 7.51 (dd, 1H); 7.74 (d, 1H); 7.82 (d, 1H); 8.22 (s, 1H).

22B 2-Cyano-6-(2-quinolinylmethoxy)naphthalene

Following the process described in example 1 (point B), starting from 2-cyano-6-hydroxynaphthalene, the title compound was prepared as a white solid with melting point 155.0-155.8°C (70% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 5.30 (s, 2H); 7.34 (d, 1H); 7.46 (dd, 1H); 7.60 (m, 2H); 7.80 (m, 5H); 8.18 (m, 2H); 8.27 (d, 1H).

22C 6-(2-Quinolinylmethoxy)-2-naphthoic acid

A solution of 35% sodium hydroxide (35 ml) was added to a solution of 2-cyano-6-(2-quinolinylmethoxy)-naphthalene (0.779, 2.5 mmol) in ethanol (130 ml) and refluxed for 24 h. After that the reaction was cooled at room temperature, added 1M HCl to pH 5 and left to stand at 5-10°C for 24 h. After this time, the resulting precipitate was filtered, washed with cold ethanol and dried over phosphorous pentoxide, thereby obtaining 0.796 g of the title compound as a white solid with

melting point 239-240°C (96% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 5.53 (s, 1H); 7.39 (dd, 1H); 7.54 (s, 1H); 7.62 (t, 2H); 7.76 (m, 3H); 8.00 (m, 4H); 8.42 (d, 1H); 8.53 (s, 1H).

5 22D Methyl 4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]carboxamidophenyl]butanoate

Following the process described in example 1 (point F), starting from 6-(2-quinolinylmethoxy)-2-naphthoic acid and methyl 4-(4-aminophenyl)butanoate, the title
10 compound was prepared (70% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.75 (q, 2H); 2.17 (t, 2H); 2.53 (t, 2H); 3.60 (s, 3H); 5.50 (s, 2H); 7.13 (d, 1H); 7.38 (dd, 1H); 7.53 (d, 1H); 7.59 (dt, 1H); 7.73 (m, 4H); 7.86 (d, 1H); 7.95 (m, 4H); 8.40 (d, 1H); 8.47
15 (s, 1H); 10.25 (s, 1H).

Example 23: 4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]carboxamidophenyl]butanoic acid

Following the process described in example 1 (point C), starting from methyl 4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]carboxamidophenyl]butanoate,
20 the title compound was prepared as a white solid with melting point 231.9-232.5°C (78% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.75 (q, 2H); 2.17 (t, 2H); 2.53 (t, 2H); 5.50 (s, 2H); 7.13 (d, 1H); 7.38 (dd, 1H); 7.53 (d, 1H); 7.59 (dt, 1H); 7.73 (m, 4H); 7.86 (d, 1H); 7.95 (m, 4H); 8.40 (d, 1H); 8.47 (s, 1H); 10.25 (s, 1H).
25

Example 24: N-[3-(1H-5-tetrazolyl)phenylmethyl]-6-(2-quinolinylmethoxy)-2-naphthalenecarboxamide

30 Following the process described in example 1 (point F), starting from 6-(2-quinolinylmethoxy)-2-naphthoic

acid and 3-(1H-5-tetrazolyl)benzylamine, the title compound was prepared as a white solid with melting point 234.5-235.0°C (68% yield).

5 ¹H N.M.R. (300 MHz, DMSO) δ ppm: 4.63 (d, 2H); 5.53 (s, 2H); 7.41 (dd, 1H); 7.56 (m, 3H); 7.63 (t, 1H); 7.75 (d, 1H); 7.81 (dt, 1H); 7.92 (m, 3H); 8.01 (m, 4H); 8.45 (d, 1H); 8.48 (s, 1H); 9.26 (t, 1H).

Example 25: methyl 4-[4-[[7-(2-quinolinylmethoxy)-2-naphthyl]carboxamidolphenyl]butanoate

10 25A 2-t-Butyldimethylsilyloxy-7-hydroxynaphthalene

t-butyldimethylsilyl chloride (4.71 g, 31.2 mmol) was added to a solution of 2,7-dihydroxynaphthalene (5 g, 0.031 mmol) and imidazole (1.9 g, 0.028 mmol) in dry DMF (30 ml), cooled at 0°C and under inert atmosphere.

15 The reaction mixture was stirred for 2.5 h. After that ethyl ether (100 ml) was added to the reaction mixture, to give a precipitate which was filtered off. The filtrate was washed with a NaCl saturated solution (3x20 ml), dried and the solvent was evaporated off, to obtain
20 a crude which was purified by flash chromatography through a silica gel column, eluting with petroleum ether:ethyl ether to obtain 6.0 g of the title compound as a white solid with melting point 104.8-105.8°C (70% yield).

25 ¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 0.30 (s, 6H); 1.07 (s, 9H); 5.86 (m, 1H); 7.00 (m, 4H); 7.69 (d, 1H); 7.70 (d, 1H).

25B: 2-t-Butyldimethylsilyloxy-7-trifluoromethylsulfoxynaphthalene

30 Pyridine (1.66 ml) and trifluoromethanesulfonic anhydride (4.32 g, 15.2 mmol) was added to a solution of

2-t-butyldimethylsilyloxy-7-hydroxynaphthalene (3.5 g, 12.7 mmol) in methylene chloride (15 ml), cooled at 0°C and under inert atmosphere. The reaction mixture was stirred at this temperature for 1 h, then diluted with ethyl ether (100 ml), washed in succession with 0.01M HCl, a NaHCO₃ saturated solution and a NaCl saturated solution, dried and the solvent was evaporated off, to obtain 5 g of the title compound as an orange oil (97% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 7.24 (m, 3H); 7.67 (s, 1H); 7.81 (d, 1H); 7.87 (d, 1H).

25C Methyl 7-t-butyldimethylsilyloxy-2-naphthalenecarboxylate

Triethylamine (3.2 ml, 23.3 mmol), dimethyl sulfoxide (31 ml) Pd(OAc)₂ (0.069 g, 0.31 mmol) and 1,3-bis(diphenylphosphino)propane (0.128 g, 0.31 mmol) were added to a solution of 2-t-butyldimethylsilyloxy-7-trifluoromethylsulfoxynaphthalene (3.6 g, 10.6 mmol) in absolute methanol (20 ml). The mixture was subjected to a carbon monoxide stream for 4 min, heated to a 75°C for 3 h under carbon monoxide atmosphere, cooled at room temperature, filtered through celite and methanol was evaporated off. The resulting solution was diluted with ethyl ether and washed in succession with water, 5% HCl, a 5% NaHCO₃ solution and a NaCl saturated solution, dried and the solvent was evaporated off, to obtain a crude which was purified by flash chromatography through a silica gel column. Eluting with petroleum ether:ethyl ether 98:2, 2.0 g of the title compound were prepared as a yellowish oil (60% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 0.27 (s, 6H); 1.04 (s,

9H); 3.95 (s, 3H); 7.20 (dd, 1H); 7.32 (d, 1H); 7.84 (m, 3H); 8.44 (s, 1H).

25D Methyl 7-hydroxy-2-naphthalenecarboxylate

5 K_2CO_3 (1.5 g, 12.5 mmol) was added to a solution of methyl 7-t-butyldimethylsilyloxy-2-naphthalene-carboxylate (1.30 g, 4.17 mmol) in tetrahydrofuran (40 ml) and methanol (40 ml) and stirred at room temperature for 4.5 h under nitrogen atmosphere. A NH_4Cl saturated solution and ethyl ether were added thereto, the phases 10 were separated and the aqueous one was extracted with ethyl ether. The combined organic extracts were washed with water, dried and the solvent was evaporated off, to obtain a crude which was purified by flash chromatography through a silica gel column. Eluting with 15 petroleum ether:ethyl acetate, 4:1, 0.96 g of the title compound were obtained (98% yield).

1H N.M.R. (300 MHz, CD_3OD) δ ppm: 3.96 (s, 3H); 7.22 (m, 2H); 7.80 (m, 3H); 8.39 (s, 1H).

20 25E Methyl 7-(2-quinolinylmethoxy)-2-naphthalenecarboxylate

Following the process described in example 1 (point B), starting from methyl 7-hydroxy-2-naphthalenecarboxylate, the title compound was prepared (97% yield).

25 1H N.M.R. (300 MHz, $CDCl_3$) δ ppm: 3.98 (s, 3H); 5.55 (s, 2H); 7.36 (d, 1H); 7.43 (dd, 1H); 7.59 (dt, 1H); 7.81 (m, 5H); 7.95 (dd, 1H); 8.16 (d, 1H); 8.23 (d, 1H); 8.47 (s, 1H).

25F 7-(2-Quinolinylmethoxy)-2-naphthoic acid

30 Following the process described in example 1 (point C), starting from methyl 7-(2-quinolinylmethoxy-

xy)-2-naphthalenecarboxylate, the title compound was prepared as a white solid with melting point 227.8-228.8°C (83% yield).

¹H N.M.R. (300 MHz, CD₃OD-CDCl₃) δ ppm: 5.52 (s, 2H);
5 7.42 (m, 2H); 7.62 (t, 1H); 7.76-7.95 (complex signal, 6H); 8.11 (d, 1H); 8.33 (d, 1H); 8.98 (s, 1H).

25G Methyl 4-[4-[[7-(2-quinolinylmethoxy)-2-naphthyl]carboxamidolphenyl]butanoate

Following the process described in example 1 (point
10 F), starting from 7-(2-quinolinylmethoxy)-2-naphthoic acid and methyl 4-(4-aminophenyl)butanoate, the title compound was prepared as a white solid with melting point 166.4-167.9°C (64% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.84 (q, 2H); 2.23 (t,
15 2H); 2.52 (t, 2H); 7.05 (d, 2H); 7.12 (d, 1H); 7.26 (dd, 1H); 7.60 (m, 8H); 8.06 (m, 4H).

Example 26: 4-[4-[[7-(2-quinolinylmethoxy)-2-naphthyl]carboxamidolphenyl]butanoic acid

Following the process described in example 1 (point
20 C), starting from methyl 4-[4-[[7-(2-quinolinylmethoxy)-2-naphthyl]carboxamidolphenyl]butanoate, the title compound was prepared as a white solid with melting point 300-302°C (80% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 1.72 (q, 2H); 1.92 (t,
25 2H); 2.50 (m, 2H); 5.54 (s, 2H); 7.15 (d, 2H); 7.45 (dd, 1H); 7.59 (d, 1H); 7.70 (m, 6H); 7.90 (m, 4H); 8.33 (s, 1H); 8.44 (m, 2H).

Example 27: N-[4-(1H-5-tetrazolyl)phenylmethyl]-7-(2-quinolinylmethoxy)-2-naphthalenecarboxamide

Following the process described in example 1 (point
30 F), starting from 7-(2-quinolinylmethoxy)-2-naphthoic

acid and 4-(1H-5-tetrazolyl)benzylamine hydrochloride, the title compound was prepared as a white solid with melting point 224.4-225.9°C (87% yield).

5 ^1H N.M.R. (300 MHz, DMSO) δ ppm: 4.58 (d, 2H); 5.53 (s, 2H); 7.44 (dd, 1H); 7.54 (m, 3H); 7.62 (t, 1H); 7.74 (d, 1H); 7.81 (m, 2H); 7.99 (m, 6H); 8.36 (s, 1H); 8.43 (d, 1H); 9.24 (t, 1H).

Example 28: ethyl 4-[2-[[7-(2-quinolinylmethoxy)-2-naphthyl]carboxamido]ethyl]benzoate

10 28A Methyl 4-cyanomethylbenzoate

Methyl 4-chloromethylbenzoate (8 g, 43.3 mmol) dissolved in ethanol (6 ml) was added to a solution of sodium cyanide (2.5 g, 51.0 mmol) in water (3 ml) and left at 100°C for 3 h. The reaction mixture was cooled at room temperature, added with ethyl ether (30 ml) and a NaCl saturated solution (10 ml). The two phases were separated and the aqueous one was extracted with ethyl ether (3x25 ml). The ether extracts were dried and the solvent was evaporated off, to obtain a crude which was purified by flash chromatography through a silica gel column. Eluting with petroleum ether:ethyl acetate, 3:2, 6.1 g of the title compound were prepared as a yellowish oil (80% yield).

20 ^1H N.M.R. (300 MHz, CDCl_3) δ ppm: 3.85 (s, 2H); 3.95 (s, 2H); 7.44 (d, 2H); 8.07 (d, 2H).

28B Methyl 4-(2-aminoethyl)benzoate

Following the process described in example 1 (point E), starting from methyl 4-cyanomethylbenzoate, the title compound was prepared as a semi-solid oil (90% yield).

30 ^1H N.M.R. (300 MHz, CD_3OD) δ ppm: 3.07 (t, 2H); 3.32 (t,

2H); 3.92 (s, 3H); 7.44 (d, 2H); 8.03 (d, 2H).

28C Ethyl 4-[2-[[7-(2-quinolinylmethoxy)-2-naphthyl]-carboxamido]ethyl]benzoate

5 Following the process described in example 1 (point F), starting from 7-(2-quinolinylmethoxy)-2-naphthoic acid and ethyl 4-(2-aminoethyl)benzoate, the title compound was prepared as a white solid with melting point 212.2-213.4°C (83% yield).

¹H N.M.R. (300 MHz, CD₃OD-CDCl₃) δ ppm: 3.02 (t, 2H);
10 3.74 (m, 2H); 3.90 (s, 3H); 5.49 (s, 2H); 6.97 (t, 1H); 7.34 (m, 3H); 7.61 (m, 2H); 7.71-7.86 (complex signal, 5H); 7.98 (d, 2H); 8.00 (m, 2H); 8.23 (d, 1H).

Example 29: 4-[2-[[7-(2-quinolinylmethoxy)-2-naphthyl]-carboxamido]ethyl]benzoic acid

15 Following the process described in example 1 (point C), starting from ethyl 4-[2-[[7-(2-quinolinylmethoxy)-2-naphthyl]carboxamido]ethyl]benzoate, the title compound was prepared as a white solid with melting point 228.0-229.0°C (63% yield).

20 ¹H N.M.R. (300 MHz, DMSO) δ ppm: 2.92 (t, 2H); 3.53 (t, 2H); 5.53 (s, 2H); 7.32 (d, 2H); 7.42 (dd, 1H); 7.51 (d, 1H); 7.63 (dt, 1H); 7.72-7.94 (complex signal, 7H); 8.01 (d, 1H); 8.06 (d, 1H); 8.25 (s, 1H); 8.44 (d, 1H); 8.70 (t, 1H).

25 Example 30: N-[4-(1H-5-tetrazolyl)phenylethyl]-7-(2-quinolinylmethoxy)-2-naphthalenecarboxamide

30A Methyl 4-[2-(t-butoxycarbonylamino)ethyl]benzoate

30 1M NaOH solution (15.4 ml) and di-t-butyl dicarbonate (2.27 g, 10.4 mmol) were added to a solution of methyl 4-(2-aminoethyl)benzoate (1.5 g, 6.96 mmol) in dioxane (30 ml) and water (15 ml) at 0°C. The reaction

mixture was stirred at room temperature for 18 h, keeping pH at 9-10 throughout the reaction by means of several additions of 1M NaOH. Dioxane was evaporated off and the aqueous residue was acidified to pH 3 with 1M HCl, extracted with ethyl acetate (3x50 ml), dried and the solvent was evaporated off to obtain 1.6 g of the title compound (92% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.42 (s, 9H); 2.83 (t, 2H); 3.38 (q, 2H); 3.88 (s, 3H); 7.27 (d, 2H); 7.98 (d, 2H).

30B 4-[2-(t-Butoxycarbonylamino)ethyl]benzoic acid

1M potassium hydroxide (34.1 ml) was added to a solution of methyl 4-[2-(t-butoxycarbonylamino)ethyl]benzoate (1.7 g, 6.82 mmol) in ethanol (380 ml) and stirred under reflux for 30 min. After that ethanol was removed, and the resulting solid residue was redissolved in water (35 ml), adjusting to pH 4-5 with 10% acetic acid, thereby obtaining a precipitate which was separated by filtration, washed with ethyl ether and dried over phosphorous pentoxide, to obtain 1.3 g of the title compound (91% yield).

¹H N.M.R. (300 MHz, CD₃OD) δ ppm: 1.41 (s, 9H); 2.82 (t, 2H); 3.30 (m, 2H); 7.32 (d, 2H); 7.92 (d, 2H).

30C 4-[2-(t-Butoxycarbonylamino)ethyl]benzamide

Triethylamine (1.7 ml, 12.34 mmol) and ethyl chloroformate (0.64 ml, 6.79 mmol) were added to a solution of 4-[2-(t-butoxycarbonylaminoethyl]benzoic acid (1.45 g, 6.17 mmol) in dry THF (100 ml). The reaction mixture was stirred at room temperature for 30 min, then subjected to an ammonia stream for 30 min., evaporated to dryness and treated with chloroform to

remove the ethyl chloroformate excess, thus obtaining 1.2 of the title compound (83% yield).

^1H N.M.R. (300 MHz, CD_3OD) δ ppm: 1.40 (s, 9H); 2.81 (t, 2H); 3.32 (m, 2H); 7.32 (d, 2H); 7.92 (d, 2H).

5 30D 4-[2-(t-Butoxycarbonylamino)ethyl]benzonitrile

A solution of 4-[2-(t-butoxycarbonylamino)ethyl]benzamide (0.5 g, 2.13 mmol) in 2 ml of DMF was added to a solution of phosphorous oxychloride (1 ml, 2.13 mmol) in dry DMF (16 ml), kept at 0°C and under inert atmosphere for 30 min. The reaction mixture was stirred at room temperature for 24 h, then poured onto ice and extracted with ethyl acetate (4x25 ml), dried and the solvent was evaporated off, to obtain 0.35 g of the title compound (77% yield).

15 ^1H N.M.R. (300 MHz, CDCl_3) δ ppm: 1.53 (s, 9H); 2.93 (t, 2H); 3.86 (t, 2H); 7.34 (d, 2H); 7.62 (d, 2H); 9.17 (s, 1H).

30E 5-[4-(2-t-Butoxycarbonylamino)phenyl]-1H-tetrazol

Following the process described in example 1 (point D), starting from 4-[2-(t-butoxycarbonylamino)-ethyl]benzonitrile, the title compound was prepared (72% yield).

^1H N.M.R. (300 MHz, CD_3OD) δ ppm: 1.37 (s, 9H); 3.30 (t, 2H); 3.50 (t, 2H); 7.39 (d, 2H); 7.88 (d, 2H).

25 30F 2-[4-(1H-5-Tetrazolyl)phenyl]ethylamine

Trifluoroacetic acid (0.37 ml, 4.8 mmol) was added to a solution of 5-[4-(2-t-butoxycarbonylamino)-phenyl]-1H-tetrazole (0.312 g, 1.20 mmol) in dry methylene chloride (4 ml) and stirred at room temperature for 18 h. After that the reaction mixture was evaporated to dryness to obtain 0.358 g of the title

compound as the trifluoroacetic acid salt (99% yield).

¹H N.M.R. (300 MHz, CD₃OD) δ ppm: 3.09 (t, 2H); 3.28 (t, 2H); 7.45 (d, 2H); 7.94 (d, 2H).

30G N-[4-(1H-5-Tetrazolyl)phenylethyl]-7-(2-quinolinyl-methoxy)-2-naphthalenecarboxamide

Following the process described in example 1 (point F), starting from 7-(2-quinolinylmethoxy)-2-naphthoic acid and 2-[4-(1H-5-tetrazolyl)phenyl]ethylamine, the title compound was prepared as a white solid with melting point 187.4-189.1°C (67% yield).

¹H N.M.R. (300 MHz, DMSO) δ ppm: 2.97 (t, 2H); 3.55 (m, 2H); 5.55 (s, 2H); 7.41-7.51 (complex signal, 4H); 7.63 (t, 1H); 7.78 (m, 3H); 7.89-8.08 (complex signal, 6H); 8.27 (s, 1H); 8.43 (s, 1H); 8.72 (t, 1H).

Example 31: ethyl 4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]methylaminocarbonyl]phenyl]butanoate

31A 2-Aminomethyl-6-methoxynaphthalene

Following the process described in example 1 (point E), starting from 2-cyano-6-methoxynaphthalene, the title compound was prepared (90% yield).

¹H N.M.R. (300 MHz, CD₃OD) δ ppm: 3.92 (s, 3H); 4.25 (s, 2H); 7.19 (dd, 1H); 7.28 (d, 1H); 7.50 (dd, 1H); 7.79-7.88 (m, 3H).

31B 2-Acetylaminomethyl-6-methoxynaphthalene

Triethylamine (2.9 ml, 20.5 mmol) and acetic anhydride (0.73 ml, 7.69 mmol) were added to a solution of 2-aminomethyl-6-methoxynaphthalene (1.23 g, 6.40 mmol) in chloroform (200 ml), cooled at -30°C. The reaction mixture was left at this temperature for 2 h, then cooled at room temperature and added with water (50 ml), the two phases were separated and the organic one

was washed with a 0.2M HCl solution, dried and the solvent was evaporated off, to obtain the title compound as a yellowish solid with melting point 163-165°C (86% yield).

5 ¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 2.03 (s, 3H); 3.90 (s, 3H); 4.53 (d, 2H); 5.95 (s, 1H); 7.13 (m, 2H); 7.35 (dd, 1H); 7.62 (s, 1H); 7.68 (m, 2H).

31C 6-Acetylaminomethyl-2-naphthol

Following the process described in example 1 (point A), starting from 2-acetylaminomethyl-6-methoxynaphthalene, the title compound was prepared as a white solid with melting point 219-222°C (93% yield).

¹H N.M.R. (300 MHz, CD₃OD) δ ppm: 2.09 (s, 3H); 4.50 (s, 2H); 7.06 (m, 2H); 7.32 (dd, 1H); 7.65 (m, 3H).

15 31D 2-Acetylaminomethyl-6-(2-quinolinylmethoxy)naphthalene

Following the process described in example 1 (point B), starting from 6-acetylaminomethyl-2-naphthol, the title compound was prepared as a semi-solid oil (65% yield).

20 ¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.93 (s, 3H); 4.49 (d, 2H); 5.44 (s, 2H); 5.85 (s, 1H); 7.25 (m, 3H); 7.51 (t, 1H); 7.67 (m, 5H); 7.78 (d, 1H); 8.07 (d, 1H); 8.15 (d, 1H).

25 31E 2-Aminomethyl-6-(2-quinolinylmethoxy)naphthalene

6M HCl (2 ml) was added to a solution of 2-acetylaminomethyl-6-(2-quinolinylmethoxy)naphthalene (0.100 g, 0.281 mmol) in dioxane (10 ml) and refluxed for 18 h. After that the reaction mixture was cooled at room temperature, diluted with water (10 ml), 1M NaOH was added to basic pH and extracted with ethyl acetate.

The organic extracts were dried and the solvent was evaporated off, to obtain 0.064 g of the title compound (73% yield).

¹H N.M.R. (300 MHz, CD₃OD) δ ppm: 3.89 (s, 2H); 5.34 (s, 2H); 7.25 (m, 2H); 7.36 (dd, 1H); 7.54 (t, 1H); 7.70 (m, 5H); 7.84 (d, 1H); 8.02 (d, 1H); 8.24 (d, 1H).

31F Ethyl 4-(4-formylphenyl)butanoate

Hexamethylenetetramine (1.6 g, 11.5 mmol) was added to a solution of ethyl 4-phenylbutanoate (2 g, 10.4 mmol) in trifluoroacetic acid (10 ml) and left at 80°C for 18 h. After that the reaction mixture was evaporated to dryness, added with a NaHCO₃ saturated solution (40 ml) and extracted with ethyl ether (4x50 ml). The ether extracts were dried and the solvent was evaporated off, to obtain a crude which was purified by flash chromatography through a silica gel column. Eluting with hexane:ethyl acetate, 9:1, 1.3 g of the title compound was prepared as a colourless oil (57% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.23 (t, 3H); 1.96 (m, 2H); 2.31 (t, 2H); 2.71 (t, 2H); 4.11 (q, 2H); 7.32 (d, 2H); 7.78 (d, 2H); 9.95 (s, 1H).

31G 4-(3-Ethoxycarbonylpropyl)benzoic acid

1 ml of Jones's reagent was added at 0°C to a solution of ethyl 4-(4-formylphenyl)butanoate (1.16 g, 5.29 mmol) in acetone (7 ml). The reaction mixture was stirred at room temperature for 18 h, then added with isopropanol (1 ml) and extracted with ethyl ether. The organic phase was dried and the solvent was evaporated off, to obtain 1.05 g of the title compound as a colourless oil (84% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.09 (t, 3H); 1.81 (m,

2H); 2.17 (t, 2H); 2.55 (t, 2H); 3.97 (q, 2H); 7.11 (d, 2H); 7.87 (d, 2H); 9.11 (m, 1H).

31H Ethyl 4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]-methylaminocarbonyl]phenyl]butanoate

5 Following the process described in example 1 (point F), starting from 2-aminomethyl-6-(2-quinolinylmethoxy)naphthalene and 4-(3-ethoxycarbonylpropyl)benzoic acid, the title compound was prepared (61% yield).

10 ¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.26 (t, 3H); 1.95 (m, 2H); 2.30 (t, 2H); 2.65 (t, 2H); 4.10 (q, 2H); 4.70 (d, 2H); 5.40 (s, 2H); 7.25 (m, 5H); 7.70 (m, 9H); 8.15 (t, 2H).

Example 32: 4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]-methylaminocarbonyl]phenyl]butanoic acid

15 Following the process described in example 1 (point C), starting from ethyl 4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]methylaminocarbonyl]phenyl]butanoate, the title compound was prepared as a white solid with melting point 185.9-187.8 (80% yield).

20 ¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.81 (t, 2H); 2.22 (t, 2H); 2.64 (t, 2H); 4.60 (d, 2H); 5.48 (m, 5H); 7.40 (m, 5H); 7.75 (m, 8H); 8.03 (m, 2H); 8.42 (d, 1H); 9.05 (t, 1H).

25 Example 33: N-[4-(1H-5-tetrazolyl)phenylpropyl]-7-(2-quinolinylmethoxy)-2-naphthalenecarboxamide

33A 3-(4-Bromophenyl)propan-1-ol

30 A solution of 4-bromocinnamic acid (5.0 g, 22 mmol) in 20 ml of dry ethyl ether was added to a suspension of aluminium lithium hydride (2.49 g, 66 mmol) in dry ethyl ether (130 ml) under inert atmosphere. The reaction mixture was stirred at room temperature for 2 hours,

then a NaCl saturated solution in water (80 ml) was slowly added, the two phases were separated and the aqueous one was extracted with ethyl acetate (3x50 ml). The organic extracts were dried and the solvent was evaporated off to obtain 3.60 g of the title compound as a yellowish oil (76% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.85 (m, 2H); 2.66 (t, 2H); 3.65 (t, 2H); 7.06 (d, 2H); 7.39 (d, 2H).

33B 3-(4-Cyanophenyl)propan-1-ol

A mixture of 3-(4-bromophenyl)propan-1-ol (2.0 g, 9.3 mmol), copper (I) cyanide (1.49 g, 16.7 mmol) and N-methylpyrrolidinone (13 ml) was stirred at 200°C for 2.5 hours. After that the reaction mixture was cooled at room temperature, poured onto a solution of diethylamine (30 g) and water (80 ml) and extracted with ethyl acetate (3x40 ml). The combined organic phases were dried and volatiles were removed, to obtain an oil from which N-methylpyrrolidinone was removed by distillation under high vacuum (0.5 torr, 85°C), thereby obtaining 0.78 g of the title compound (52% yield).

¹H N.M.R. (300 MHz, CDCl₃) δ ppm: 1.85 (m, 2H); 2.42 (s broad, 1H); 2.76 (t, 2H); 3.64 (t, 2H); 7.29 (d, 2H); 7.53 (d, 2H).

33C 3-(4-Cyanophenyl)propyl methanesulfonate

Triethylamine (0.54 ml, 4.03 mmol) and methanesulfonyl chloride (0.30 ml, 4.03 mmol) were added to a solution of 3-(4-cyanophenyl)propan-1-ol (0.50 g, 3.10 mmol) in dry methylene chloride (15 ml), cooled at 0°C and under inert atmosphere. The reaction mixture was stirred at 0°C for 2 hours, after that was diluted with methylene chloride (50 ml), washed in succession with

0.05M HCl, with a NaCl saturated solution, dried and the solvent was evaporated off. 0.675 g of the title compound were obtained as a semi-solid oil (94% yield).

5 ^1H N.M.R. (300 MHz, CDCl_3) δ ppm: 1.89 (m, 2H); 2.63 (t, 2H); 2.85 (s, 3H); 4.05 (t, 2H); 7.14 (d, 2H); 7.38 (d, 2H).

33D 4-(1H-5-Tetrazolyl)azidopropylbenzene

10 Following the process described in example 1 (point D), starting from 3-(4-cyanophenyl)propyl methanesulfonate, the title compound was prepared (70% yield).

^1H N.M.R. (300 MHz, CDCl_3) δ ppm: 1.79 (m, 2H); 2.63 (t, 2H); 3.18 (t, 2H); 7.21 (d, 2H); 7.98 (d, 2H).

15 33E 3-[4-(1H-5-Tetrazolyl)phenyl]propylamine hydrochloride

20 Following the process described in example 1 (point E), starting from 4-(1H-5-tetrazolyl)azidopropylbenzene, the title compound was prepared as a white solid (crystallized from methanol) which decomposes at 251°C (87% yield).

^1H N.M.R. (300 MHz, CD_3OD) δ ppm: 2.03 (m, 2H); 2.83 (t, 2H); 3.00 (t, 2H); 7.48 (d, 2H); 7.97 (d, 2H).

33F N-[4-(1H-5-Tetrazolyl)phenylpropyl]-7-(2-quinolinylmethoxy)-2-naphthalenecarboxamide

25 Following the process described in example 1 (point F), starting from 7-(2-quinolinylmethoxy)-2-naphthoic acid and 3-[4-(1H-5-tetrazolyl)phenyl]propylamine, the title compound was prepared as a white solid with melting point 218.0-219.8°C (65% yield).

30 ^1H N.M.R. (300 MHz, CD_3OD) δ ppm: 1.91 (m, 2H); 2.73 (t, 2H); 3.33 (t, 2H); 5.52 (s, 2H); 7.42 (dd, 1H), 7.45-

7.54 (complex signal, 3H); 7.63 (t, 1H); 7.72-7.84 (complex signal, 3H); 7.88-8.00 (complex signal, 4H); 8.03 (dd, 1H); 8.25 (d, 1H); 8.28 (s, 1H); 8.44 (d, 1H); 8.63 (t broad, 1H).

5 Biological activity tests

The antagonistic activity on LTD₄ of the compounds of the present invention is determined by means of an inhibition test of the [³H]-LTD₄ receptor binding in guinea-pig lung membranes, and a test of inhibition of LTD₄-induced contractions in the mienteric plexus of guinea-pig isolated ileum.

10 [³H]-LTD₄ receptor binding inhibition test in guinea-pig lung membranes

Guinea pig lung membranes, containing the LTD₄ receptors, are purified following the method described by Mong et. al. (Mong et al., Prostaglandins, 28, 805 (1984)). These purified membranes (150 µg/ml) are added to an incubation mixture containing 10 mM of PIPES buffer (piperazin-N,N'-bis(2-ethanesulfonic acid) (pH 7.4), 10 mM of CaCl₂, 10 mM of 5 MgCl₂, 2 mM of cysteine, 2 mM of glycine, 0.5 nM of [³H]-LTD₄ (4700-6400 GBq/mmol) and different concentrations of the product under test in a final volume of 310 µl. The reaction mixture is incubated for 30 minutes at 25°C.

25 The radioligand bound to the membranes is separated from the free one by dilution with 4 ml washing buffer (10 mM Tris-HCl (pH 7.4) and 100 mM NaCl) at 0°C and filtration with Whatman GF/B filters, by means of a Brandel Cell Harvester. The filters are washed 4 times with a total volume of 16 ml of washing buffer at 0°C. The radioactivity present in the filters is determined

by liquid scintillation.

The specific binding is defined as the difference between the total binding of [³H]-LTD₄ and the non-specific binding determined in the presence of 1 μM LTD₄. The data obtained in the competition tests are analyzed by a computational program, which determines the inhibition constant of each compound (K_i) by means of the Cheng-Prusoff equation (Cheng et al., Biochem. Pharmacol., 22, 3094 (1973)).

$$K_i = IC_{50} / (1 + [L] / K_d)$$

wherein IC₅₀ is the concentration of compound which displaces a 50% of the bound radioligand, [L] is the concentration of [³H]LTD₄ free in the test and K_d is the dissociation constant of the LTD₄ obtained in an independent way by means of Scatchard analysis.

Table 1 shows some of the activity values found for the compounds of the present invention.

Inhibition test of the contractions induced by LTD₄ in the mienteric plexus of guinea-pig isolated ileum.

The antagonistic activity of the compounds of the present invention in the isolated organ was evaluated as its ability to inhibit the contraction caused by LTD₄ in the mienteric plexus of the ileum of Dunkin Hartley male albino guinea-pig, weighing 300-350 g (Cristol J.P. and Sirois P. Res. Commun. Chem. Pathol., 52, 423 (1988)).

The smooth muscle of guinea-pig ileum exhibits sensitivity to leukotrienes and especially to LTD₄, which acts as primary mediator in the inflammatory and allergic response (Carnathan G.W. et al. Agents Actions, 20, 124 (1987)).

The mienteric plexus is extracted from a 2-3 cm

segment of the terminal portion of the guinea-pig ileum, previously sacrificed by cervical dislocation. The plexus is put, at a tension of 0.5 g, in a 5 ml organ bath, containing a solution of Tyrode (137 mM NaCl, 2.7 mM KCl, 1.4 mM CaCl_2 , 0.4 mM NaH_2PO_4 , 11.9 mM NaHCO_3 , 0.8 mM MgSO_4 , 5.5 mM glucose), saturated with carbogen gas (95% O_2 -5% CO_2) at 37°C. The solution also contains indomethacin (3.3 μM) and atropine (0.4 μM) to remove the action of the intrinsic prostaglandins and the cholinergic responses.

After a 45 minute stabilization period a maximum isotonic response is obtained (100% contractile response) adding to the bath chamber the LTD_4 agonist (3 nM). This process is repeated until the same contraction response is obtained twice. The isometric measures are made in an isotonic transducer.

After stabilization is restored, the product under test is incubated at different concentrations (dissolved in 0.1% final concentration DMSO) for 2.5 minutes, and after that the contraction with LTD_4 is induced again.

The antagonistic activity is expressed as IC_{50} , the concentration of compound which reduces by 50% the maximum contraction.

Table 1 shows some of the values of activity found for the compounds of the present invention.


63

Table 1

Compound Example N°.	Inhibition of the [³ H]-LTD ₄ receptor binding K _i (nM)	Inhibition of contractions induced in the ileum by LTD ₄ IC ₅₀ (nM)
5		
	1	9.2±2
	2	29.5±3
	3	12.0±3
10	5	34.0±5
	6	32.7±3
	10	38.5±6.3
	14	68.0±5
	18	134±18
15	20	190±21
	21	5.5±0.5
	23	30.0±1.2
	24	28.0±3
	26	1.5±0.2
20	27	7.8±2
	29	1.0±0.1
	30	3.4±1.9
	32	44±15
	33	13.1±2
25		

CLAIMS

1. A compound of formula I,



10 wherein:

the substituent containing A is bound to the 6- or 7-
position of the 2-naphthol system;

the substituent containing B is bound to the benzene
ring at any free position;

15 -R¹ is hydrogen or methyl;

-R² is hydrogen, fluorine, chlorine or -OCH₃, which is
bound to the naphthalene system at any positions except
the 2- and the one occupied by the other substituent;

-R³ is hydrogen, fluorine, chlorine or bromine;

20 -A- is a -CO-NR⁴- or -NR⁴-CO- group, wherein R⁴ is
hydrogen or methyl;

-B is a 5-tetrazolyl or -COOR⁵ group, wherein R⁵ is
hydrogen, a (C₁-C₄)-alkyl or a phenylalkyl group of less
than 10 carbon atoms;

25 m is 0 or 1;

n and p are integers from 0 to 6, with the proviso that
n + p is less or equal to 6;

as well as the solvates and pharmaceutically acceptable
salts thereof.

30 2. A compound according to claim 1, wherein R₂ is
hydrogen, B is a 5-tetrazolyl or COOR⁵ group, and R⁵ is

hydrogen, methyl, ethyl or benzyl.

3. A compound according to claim 1 or 2, wherein R^3 is hydrogen or chlorine, and -A- is -CONH- or -NHCO-.

4. A compound according to any one of claims 1 to 3,
5 wherein the substituent containing A is bound to the 6-position of the 2-naphthol system.

5. A compound according to claim 4, wherein R^1 is hydrogen, m is 1, and -A- is -NHCO-.

6. A compound according to claim 4, wherein -A- is
10 -CONH-.

7. A compound according to claim 6, wherein n and p are integers from 0 to 3.

8. A compound according to any one of claims 1 to 3,
15 wherein the substituent containing A is bound to the 7-position of the 2-naphthol system.

9. A compound according to claim 8, wherein R^1 is hydrogen, m is 1 and -A- is -CONH-.

10. A compound according to claim 8, wherein m is 0 and A is -CONH-.

11. A compound according to claim 10, wherein n and p
20 are integers from 0 to 3.

12. A compound according to claim 1 selected from the following ones:

25 N-[4-(1H-5-tetrazolyl)phenylmethyl]-2-[6-(2-quinolinyl-methoxy)-2-naphthyl]propanamide;

N-[3-(1H-5-tetrazolyl)phenylmethyl]-2-[6-(2-quinolinyl-methoxy)-2-naphthyl]propanamide;

N-[2-(1H-5-tetrazolyl)phenylmethyl]-2-[6-(2-quinolinyl-methoxy)-2-naphthyl]propanamide;

30 N-[4-(1H-5-tetrazolyl)methylphenyl]-2-[6-(2-quinolinyl-methoxy)-2-naphthyl]propanamide;


- N-[4-(1H-5-tetrazolyl)methylphenyl]-2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamide (sodium salt);
4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamido]-benzoic acid;
- 5 4-[4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamido]phenyl]butanoic acid;
4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidomethyl]benzoic acid;
3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamidomethyl]benzoic acid;
- 10 4-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamido]-phenylacetic acid;
3-[2-[6-(2-quinolinylmethoxy)-2-naphthyl]propanamido]-phenylacetic acid;
- 15 4-[4-[2-[6-[(7-chloro-2-quinolinyl)methoxy]-2-naphthyl]propanamido]phenyl]butanoic acid;
N-[4-(1H-5-tetrazolyl)phenylmethyl]-6-(2-quinolinylmethoxy)-2-naphthaleneacetamide;
4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]carboxamido]-phenyl]butanoic acid;
- 20 N-[3-(1H-5-tetrazolyl)phenylmethyl]-6-(2-quinolinylmethoxy)-2-naphthalenecarboxamide;
4-[4-[[7-(2-quinolinylmethoxy)-2-naphthyl]carboxamido]-phenyl]butanoic acid;
- 25 N-[4-(1H-5-tetrazolyl)phenylmethyl]-7-(2-quinolinylmethoxy)-2-naphthalenecarboxamide;
4-[2-[[7-(2-quinolinylmethoxy)-2-naphthyl]carboxamido]-ethyl]benzoic acid;
- 30 N-[4-(1H-5-tetrazolyl)phenylethyl]-7-(2-quinolinylmethoxy)-2-naphthalenecarboxamide;
4-[4-[[6-(2-quinolinylmethoxy)-2-naphthyl]methylamino-

carbonyl]phenyl]butanoic acid;


N-[4-(1H-5-tetrazolyl)phenylpropyl]-7-(2-quinolinylmethoxy)-2-naphthalenecarboxamide.

13. A process for the preparation of the compounds of general formula I of claim 1, and of the pharmaceutically acceptable salts thereof, wherein:


a) when in formula I -A- is $-\text{CO}-\text{NR}^4-$, then a compound of general formula II,



wherein R^1 , R^2 , R^3 and m have the above defined meanings, is reacted with a compound III,



20 wherein R^4 , n and p have the above defined meanings and D can be equivalent to the group B in I or, when B in formula I is COOH , then D contains suitable carboxy-protecting group; the reaction between II and III being carried out in the presence of a carboxy-activating agent and a base, to obtain a compound of formula IVa,




which is equivalent to I, or is converted into I by

removing of the carboxy-protecting group;


b) when in formula I -A- is $-NR^4-CO-$, then a compound of general formula V,

5



10


wherein R^1 , R^2 , R^3 , R^4 and m have the above defined meanings, is reacted with a compound VI



15

wherein n , p and D have the above defined meanings; the reaction between V and VI being carried out in the presence of a carboxy-activating agent and a base, to obtain a compound of formula IVb,

20



25

which coincides with I, or is converted into I as described above for the compound IVa;

c) and, if desired, the compound of general formula I is converted into the desired salt, by treatment with a base or a suitable ion-exchanger, according to conventional methods.

30

14. The use of a compound of any one of claims 1 to 12 in the preparation of a medicament for the therapeutical

treatment of leukotriene-mediated diseases.

15. The use according to claim 14, wherein the leukotriene-mediated diseases are of inflammatory or allergic type.

5 16. The use according to claim 15, wherein the inflammatory or allergic diseases are: bronchial asthma, allergic rhinitis, allergic conjunctivitis, rheumatoid arthritis, osteoarthritis, tendinitis, bursitis or psoriasis.

10 17. The use according to claim 14, wherein the leukotriene-mediated diseases are of cardiovascular type.

15 18. The use according to claim 17, wherein the diseases of cardiovascular type are: cardiac ischemia, cardiac infarction, coronary spasm, cardiac anaphylaxis, cerebral oedema or endotoxic shock.

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/EP 95/02970

 A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07D401/12 A61K31/47 C07D215/14 C07D215/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 348 155 (RORER INTERNATIONAL (OVERSEAS) INC.) 27 December 1989 see claims	1, 14
A	EP,A,0 315 399 (RORER INTERNATIONAL (OVERSEAS) INC.) 10 May 1989 see claims	1, 14
A	JOURNAL OF MEDICINAL CHEMISTRY, vol. 35, no. 14, WASHINGTON US, pages 2501-2524, JOHN H. MUSSER ET AL. '5-Lipoxygenase: properties, ...' * page 2514, figure 10, column on the right *	1, 14



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *A* document member of the same patent family

Date of the actual completion of the international search

16 November 1995

Date of mailing of the international search report

22. 11. 95

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+ 31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/EP 95/02970

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,5 084 575 (AMERICAN HOME PRODUCTS CORPORATION) 28 January 1992 cited in the application see claims -----	1,14
A	US,A,4 579 866 (USV PHARMACEUTICAL CORP.) 1 April 1986 cited in the application see claims -----	1,14

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No.

PCT/EP 95/02970

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-348155	27-12-89	US-A- 4920131 WO-A- 8912628	24-04-90 28-12-89
EP-A-315399	10-05-89	US-A- 4920132 AU-B- 633475 AU-B- 2794689 JP-T- 3500889 WO-A- 8904305 US-A- 4920131 US-A- 5059610	24-04-90 04-02-93 01-06-89 28-02-91 18-05-89 24-04-90 22-10-91
US-A-5084575	28-01-92	US-A- 4960892 AU-B- 654292 AU-B- 8617191 CA-A- 2089262 EP-A- 0547148 JP-T- 6500997 WO-A- 9204325 US-A- 5250693 US-A- 5208344 AU-B- 611699 AU-B- 1920988 CA-A- 1330999 EP-A, B 0301813 GB-A, B 2207428 IE-B- 60240 JP-A- 1100144 PT-B- 88116 CA-A- 1331000 AU-B- 629868 AU-B- 4436189 CA-A- 1331001 DE-D- 68914646 DE-T- 68914646 DK-B- 169545 EP-A- 0396839 ES-T- 2063143 GB-A, B 2231570 IE-B- 61967 JP-A- 2311463	02-10-90 03-11-94 30-03-92 07-03-92 23-06-93 27-01-94 19-03-92 05-10-93 04-05-93 20-06-91 02-02-89 26-07-94 01-02-89 01-02-89 15-06-94 18-04-89 01-03-95 26-07-94 15-10-92 15-11-90 26-07-94 19-05-94 21-07-94 28-11-94 14-11-90 01-01-95 21-11-90 30-11-94 27-12-90

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 95/02970

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4579866	01-04-86	AU-B- 4311685	05-12-85
		EP-A- 0178381	23-04-86
		JP-A- 61000054	06-01-86
<hr/>			

THIS PAGE BLANK (USPTO)